TRAILS around HEADQUARTERS



Rlue-2

Riparian zones

Riparian zones occur along creek and river margins and often contain characteristic vegetation such as river birch, sycamore and silver maple. Because of their proximity to water, these areas serve as habitat for frogs, egg laying sites for salamanders, and watering holes for other wildlife including birds. To see one of our best examples, hike the Pea Ridge loop trail and look for areas that fit this description. The Acadian flycatcher is just one species that often uses these areas.

Grassland

Once common to the Bluegrass Region of Kentucky, grasslands were integral for wildlife and the early people of the state. Native grasslands consist of tall grasses and wildflowers which are ideal for several wildlife and bird species. Grasslands provide bird species with a wide array of seeds and insects for feeding. Salato's grasslands can be seen as you pull onto Sportman's Lane and on Salato's Prairie Trail. During migration and the breeding season, prairie warbler and indigo bunting can be found in brushier areas. Year-round, field sparrows, song sparrows and bluebirds are easily found.

Lakes


Lakes are formed when water gathers in an area where the soil is impenetrable. The Upper and Lower Sportsman Lakes are man-made, and managed with waterfowl and fish in mind. In winter, a diverse array of waterfowl can be observed, attracted by the variety of food and cover located in and around the lake. The woods surrounding the lakes are usually filled with migrant warblers during spring and fall. Throughout the summer look for orioles and their unique pouch-like nests hanging from limbs of large shade trees surrounding the lakes. Eastern phoebes and swallows catch insects over the water during the breeding season. Keep an eye out for goslings in early May as the Canada geese nest right here at the lakes.

Moist soil areas

Water levels in ponds and wetlands naturally rise and fall on a seasonal basis. When biologists attempt to mimic this natural system it is called moist soil management. Lowering water levels during the summer months encourages vegetation to grow. Shorebirds will frequent the draining areas in the late summer and fall. During the fall and winter, the flooded vegetation in larger moist soil units can provide food and cover for migrating and wintering waterfowl. Biologists can target species groups by simply altering water levels. Common yellowthroat can be observed during migration and the breeding season. Sandpipers feed along the shoreline during migration.

Woodland

This type of habitat is great for many species of birds, as the tree canopy overhead provides excellent hiding places. On the Pea Ridge and Habitrek trails you'll be able to experience a central Kentucky oak-hickory forest. Mostly composed of oak and hickory tree species, this type of habitat is also known for the Kentucky coffee tree and rock elm. Oak-hickory forests lack the characteristic species found farther east in the state, such as eastern hemlock, magnolias and yellow buckeye. Throughout the year, you can see Carolina chickadee, downy woodpecker, white-breasted nuthatch and tufted titmouse. Various warbler species may be found in the woodlands during spring and fall migration. When the breeding season arrives, the red-eyed vireo and eastern wood pewee are easily heard.



Road

Habitrek Trail

----- Salato exhibit trails

- - Pea Ridge Trail

Prairie Trail

Salato exhibit path

Warbler Ridge Trail

Property boundary

Picnic shelter

Trail marker

🛉 🛉 Restroom

Bench

Gate

1#



Kentucky Dept. of Fish & Wildlife Resources 1 Sportsman's Lane, Frankfort, KY 40601 1-800-858-1549 • *fw.ky.gov*



BIRD SPECIES at KENTUCKY FISH and WILDLIFE HEADQUARTERS



Date

Observers

Salato, Lakes and Pea Ridge Trail

LEGEND

- 4 = Seen or heard most visits 3 = Common
- 2 = Uncommon 1 = Rare
- . = Unlikely in that season
- Sp = SprinqSu = Summer Fa = Fall Wi = Winter



| GEESE and DUCKS | Sn | Su | Fa | Wi |
|---|--------|--------|--------|--------|
| Snow Goose | Ъþ | | Iu | 1 |
| Ross's Goose | | • | • | 1 |
| Canada Goose | 4 | 4 | 4 | 4 |
| Wood Duck | | 3 | | |
| American Widgeon | | | | |
| American Black Duck | | | | 1 |
| Mallard | 4 | 4 | 4 | 4 |
| Blue-winged Teal | 1 | | • | |
| Northern Shoveler | | | | 2 |
| Northern Pintail | | | | 1 |
| Green-winged Teal | | | | |
| Canvasback | | | | 1 |
| Redhead | | | | _ |
| Ring-necked Duck | • | | | 1 |
| Lesser Scaup | • | | 1 | |
| Bufflehead | • | | | 1 |
| Common Goldeneye | • | | • | |
| Common Merganser | • | | | 1 |
| Ruddy Duck | • | • | • | 1 |
| Ruddy Duck | • | • | • | 1 |
| QUAIL and TURKEY Northern Bobwhite Wild Turkey | 1 | 1 | 1 | 1 |
| GREBES Pied-billed Grebe | 2 | 1 | 2 | 1 |
| CORMORANTS Double-crested Cormorant | • | 1 | | |
| HERONS Great Blue Heron Green Heron | 3 1 | 3 3 | | 3 |
| VULTURES Turkey Vulture Black Vulture | 4 3 | 4 3 | 4 3 | 4 3 |
| HAWKS, EAGLES and FALCONS Osprey | 1 | | 1 | |
| Bald Eagle | т | • | т | · |
| Northern Harrier | • | · | · | • |
| | • | • | • | • |

| Northern Harrier | | | |
|---------------------|---|---|---|
| Sharp-shinned Hawk | 1 | 1 | 1 |
| Cooper's Hawk | 1 | 1 | 2 |
| Red-shouldered Hawk | | | • |
| Broad-winged Hawk | | | |
| Red-tailed Hawk | 3 | 2 | 2 |
| American Kestrel | | | |
| Merlin | | | |
| | | | |
| COOT and CRANE | | | |

American Coot Sandhill Crane

| PLOVERS and SANDPIPERS |
|---|
| Killdeer |
| Spotted Sandpiper |
| Solitary Sandpiper |
| Lesser Yellowlegs |
| American Woodcock |
| GULLS and TERNS |
| Ring-billed Gull |
| DOVEC |
| DOVES |
| Rock Pigeon |
| Mourning Dove |
| CUCKOOC |
| CUCKOOS |
| Yellow-billed Cuckoo Black-billed Cuckoo |
| |
| OWI C |

OWLS

Eastern Screech Owl Great Horned Owl Barred Owl

NIGHTJARS Common Nighthawk -widow

| T | Chuck-will's- |
|---|---------------|
| | |

SWIFTS and HUMMINGBIRDS

232. Chimney Swift Ruby-throated Humm. 232.

Sp Su Fa Wi

2 3 2 2

2 1 . .

1 1 . .

. . . .

2 1 2 .

1 . . .

1 1 1 1

4 4 4 4

2 3 1 .

1 . . .

1 1 1 1

1 1 1 1

1 1 1 1

. 1 1 .

. . . .

KINGFISHERS

Belted Kingfisher 3 3 3 3

WOODPECKERS

Red-bellied Woodpecker 4 4 4 4 Yellow-bellied Sapsucker 2 . . 1 Downy Woodpecker 4 4 4 4 Hairy Woodpecker 2 2 2 2 Northern Flicker 3 3 3 3 Pileated Woodpecker 2 2 2 2

FLYCATCHERS

2

1

2

3 1 . 2

. . 1 1

| Eastern Wood Pewee | 4 | 4 | 1 | |
|---------------------------|---|---|---|---|
| Yellow-bellied Flycatcher | | | | |
| Acadian Flycatcher | | 2 | 1 | |
| Least Flycatcher | 1 | | 1 | |
| Eastern Phoebe | 2 | 3 | 2 | 1 |
| Great-crested Flycatcher | 3 | 2 | | |
| Eastern Kingbird | 3 | 4 | | |

VIREOS White-

| e-eyed | Vireo | 2 | 3 | 1 | |
|--------|-------|---|---|---|--|
| | | | | | |

| | Sp | Su | Fa | Wi |
|-------------------------|----|----|----|----|
| Yellow-throated Vireo | 1 | | | |
| Blue-headed Vireo | 1 | • | • | • |
| Warbling Vireo | 2 | 3 | 1 | |
| Philadelphia Vireo | • | | • | • |
| Red-eyed Vireo | 2 | 3 | 1 | • |
| | | | | |
| JAYS and CROWS | | | | |
| Blue Jay | 4 | 4 | 4 | 4 |
| American Crow | 4 | 4 | 4 | 4 |
| | | | | |
| SWALLOWS | | | | |
| Purple Martin | 2 | 3 | • | • |
| Tree Swallow | | 3 | | • |
| N. Rough-winged Swallow | 1 | 1 | • | • |
| Barn Swallow | 3 | 4 | 1 | • |
| | | | | |
| CHICKADEES and TITMI | - | | | |
| Carolina Chickadee | | 4 | 4 | 4 |
| Tufted Titmouse | 4 | 4 | 4 | 4 |
| | | | | |
| NUTHATCHES and | | | | |
| CREEPERS | , | , | , | , |

| White-breasted Nuthatch | 4 | 4 | 4 | 4 |
|-----------------------------------|---|---|---|---|
| Red-breasted Nuthatch | • | • | | • |
| Brown Creeper | 1 | | | 1 |
| | | | | |
| WRENS | | | | |
| Carolina Wren | 4 | 4 | 4 | 4 |
| House Wren | 1 | 3 | 1 | |
| Winter Wren | | | | 1 |
| | | | | |
| THRUSHES and ALLIES | | | | |
| Golden-crowned Kinglet | 1 | | 2 | 2 |
| Ruby-crowned Kinglet | 1 | | 1 | 1 |
| Blue-Gray Gnatcatcher | 2 | 2 | 1 | |

4 4 4 3

343.

232.

4 4 4 4

1 . . .

| Blue-Gray Gnatcatcher | 2 | 2 | 1 | | |
|-----------------------|---|---|---|---|--|
| Eastern Bluebird | 3 | 2 | 2 | 3 | |
| Gray-cheeked Thrush | 1 | | 1 | | |
| Swainson's Thrush | 1 | | 1 | | |
| Hermit Thrush | 1 | | 1 | 1 | |
| Wood Thrush | 2 | 2 | • | | |
| American Robin | 4 | 4 | 4 | 4 | |
| | | | | | |

MIMICS

Northern Mockingbird Gray Catbird Brown Thrasher

STARLINGS

| European | Starling |
|----------|----------|
| | |

PIPITS American Pipit

WAXWINGS

Cedar Waxwing

WOOD WARBI

NEW WORLD SPARROWS

TANAGERS

Scarlet Tanager

Summer Tanager

and ALLIES Eastern Towhee American Tree Spa **Chipping Sparrow** Field Sparrow Fox Sparrow Song Sparrow Swamp Sparrow White-throated Sp White-crowned Sp Dark-eyed Junco

| WAXWINGS | Sp | Su | Fa | Wi | CARDINALS and | Sp | Su | Fa | Wi |
|--------------------------|----|----|----|----|-----------------------------|----|----|----|----|
| Cedar Waxwing | 2 | 2 | 2 | 1 | ALLIES | • | | | |
| | | | | | Northern Cardinal | 4 | 4 | 4 | 4 |
| WOOD WARBLERS | | | | | Rose-breasted Grosbeak | 1 | | 1 | |
| Blue-Winged Warbler | | | | | Blue Grosbeak | 1 | | | |
| Golden-Winged Warbler | | | | | Indigo Bunting | 3 | 4 | 2 | • |
| Tennessee Warbler | | | 1 | | | | | | |
| Orange-crowned Warbler | | | | | BLACKBIRDS and | | | | |
| Nashville Warbler | | | | | ORIOLES | | | | |
| Northern Parula | 2 | 2 | | | Bobolink | | | | |
| Yellow Warbler | 2 | 1 | | | Red-winged Blackbird | 4 | 4 | 4 | 1 |
| Chestnut-sided Warbler | 1 | | 1 | | Eastern Meadowlark | 3 | 3 | 1 | 1 |
| Magnolia Warbler | 1 | | 2 | | Common Grackle | 4 | 4 | 3 | 1 |
| Cape May Warbler | 1 | | 1 | | Brown-headed Cowbird | 4 | 3 | 1 | 1 |
| Black-throated Blue War | | | | | Rusty Blackbird | | | | |
| Yellow-rumped Warbler | 2 | | 2 | 2 | Orchard Oriole | 2 | 2 | 1 | |
| Black-throated Green War | | | 1 | | Baltimore Oriole | 2 | 4 | 1 | |
| Blackburnian Warbler | 1 | | 1 | | | | | | |
| Yellow-throated Warbler | | | 1 | | FINCHES | | | | |
| Pine Warbler | 1 | | | | American Goldfinch | 4 | 4 | 4 | 4 |
| Prairie Warbler | 1 | 2 | 1 | | Purple Finch | | | | |
| Palm Warbler | 1 | | 1 | | | 3 | 2 | 3 | 2 |
| Bay-breasted Warbler | | | | | House Sparrow | 4 | 4 | 4 | 4 |
| Blackpoll Warbler | 1 | | | | · | | | | |
| Black-and-white Warbler | 1 | | 2 | | A total of 175 species have | be | en | | |
| American Redstart | 1 | | 2 | | recorded here. | | | | |
| Prothonotary Warbler | | | | | | | | | |
| Worm-eating Warbler | | | | | | | | | |
| Ovenbird | 3 | 2 | 1 | | ADDITIONAL SPECIES | | | | |
| Northern Waterthrush | 1 | | | | | | | | |
| Louisiana Waterthrush | 1 | 1 | | | | | | | |
| Kentucky Warbler | 2 | 3 | 1 | | | | | | |
| Connecticut Warbler | | | | | | | | | |
| Mourning Warbler | 1 | | 1 | | | | | | |
| Common Yellowthroat | 2 | 3 | 2 | | | | | | |
| Hooded Warbler | | | | | | | | | |
| Wilson's Warbler | | | | | | | | | |
| Canada Warbler | 1 | | 1 | | | | | | |
| Yellow-breasted Chat | 1 | 2 | 1 | | | | | | |
| 74146506 | | | | | | | | | |

| | 4 | 4 | 4 | 3 | |
|--------|---|---|---|---|--|
| arrow | | | | 1 | |
| | 4 | 3 | 2 | | |
| | 4 | 4 | 2 | 1 | |
| | 1 | | 1 | 2 | |
| | 4 | 4 | 4 | 4 | |
| | 1 | | 2 | 1 | |
| parrow | 3 | | 2 | 4 | |
| barrow | | | | 1 | |
| | 3 | | 1 | 4 | |

1 1 1 .

1 3 1 .